arXiv:math/0112186v2 [math.AC] 25 Feb 2003

THE HILBERT FUNCTIONS OF ACM SETS OF POINTS IN $\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$

ADAM VAN TUYL

Abstract. If $\mathbb{X}$ is a set of points in $\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$, then the associated coordinate ring $R/I_{\mathbb{X}}$ is an $\mathbb{N}^k$-graded ring. The Hilbert function of $\mathbb{X}$, defined by $H_{\mathbb{X}}(\underline{i}) := \dim_{\mathbf{k}}(R/I_{\mathbb{X}})_{\underline{i}}$ for all $\underline{i} \in \mathbb{N}^k$, is studied. Since the ring $R/I_{\mathbb{X}}$ may or may not be Cohen-Macaulay, we consider only those $\mathbb{X}$ that are ACM. Generalizing the case of $k = 1$ to all k, we show that a function is the Hilbert function of an ACM set of points if and only if its first difference function is the Hilbert function of a multi-graded artinian quotient. We also give a new characterization of ACM sets of points in $\mathbb{P}^1 \times \mathbb{P}^1$, and show how the graded Betti numbers (and hence, Hilbert function) of ACM sets of points in this space can be obtained solely through combinatorial means.

1. Introduction

Let $R = \mathbf{k}[x_{1,0}, \ldots, x_{1,n_1}, \ldots, x_{k,0}, \ldots, x_{k,n_k}]$ with $\deg x_{i,j} = e_i$ be the $\mathbb{N}^k$-graded coordinate ring associated to $\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$. A point $P = \mathcal{P}_1 \times \cdots \times \mathcal{P}_k \in \mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$, with $\mathcal{P}_i \in \mathbb{P}^{n_i}$, corresponds to a prime $\mathbb{N}^k$-homogeneous ideal I_P of height $\sum_{i=1}^{k} n_i$ in R. Furthermore, $I_P = (L_{1,1}, \ldots, L_{1,n_1}, \ldots, L_{k,1}, \ldots, L_{k,n_k})$ where $\deg L_{i,j} = e_i$ and $(L_{i,1}, \ldots, L_{i,n_i})$ is the defining ideal of $\mathcal{P}_i \in \mathbb{P}^{n_i}$. If $\mathbb{X} = \{P_1, \ldots, P_s\} \subseteq \mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$, then $I_{\mathbb{X}} = \bigcap_{i=1}^{s} I_{P_i}$, where I_{P_i} corresponds to P_i, is the $\mathbb{N}^k$-homogeneous ideal of R associated to $\mathbb{X}$. The ring $R/I_{\mathbb{X}}$ inherits an $\mathbb{N}^k$-graded structure. The Hilbert function of $\mathbb{X}$ is then defined by $H_{\mathbb{X}}(\underline{i}) = \dim_{\mathbf{k}}(R/I_{\mathbb{X}})_{\underline{i}}$ for all $\underline{i} \in \mathbb{N}^k$. In this paper we study these Hilbert functions, thereby building upon [5, 13].

Each ideal I_{P_i} is also homogeneous with respect to the standard grading, so I_{P_i} defines a linear variety of dimension $k-1$ in $\mathbb{P}^{N-1}$ where $N = \sum_{i=1}^{k}(n_i + 1)$. One can therefore take the point of view that our investigation of sets of points in $\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$ is an investigation of reduced unions of linear varieties with extra hypotheses on the generators to ensure $I_{\mathbb{X}}$ is $\mathbb{N}^k$-homogeneous.

Ideally, we would like to classify those functions that arise as the Hilbert function of a set of points in $\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$. However, besides the case $k = 1$ which is dealt with in [3] and [4], such a classification continues to elude us. Though some properties of the Hilbert function are known if $k > 1$ (cf. [5, 13]), even for sets of points in $\mathbb{P}^1 \times \mathbb{P}^1$ this problem remains open.

The proof of the characterization for the case $k = 1$ relies, in part, on the fact that the coordinate ring of any finite set of points in $\mathbb{P}^n$ is always Cohen-Macaulay (CM). However, if $k > 1$, we show how to construct sets of points which fail to be CM. In fact, for each integer $l \in \{1, \ldots, k\}$, we can construct a set of points with $\operatorname{depth} R/I_{\mathbb{X}} = l$. The failure of $R/I_{\mathbb{X}}$ to be CM in general provides an obstruction to generalizing the proofs of [3, 4].

We therefore restrict our investigation to sets of points that arithmetically Cohen-Macaulay (ACM). With this extra hypothesis on our set of points, we can generalize the proof for the case $k = 1$ as given in [3] to all k. In particular, we show that $H_{\mathbb{X}}$ is the Hilbert function of an

---

Key words and phrases. Hilbert function, points, multi-projective space, Cohen-Macaulay rings, partitions.
Updated: Feb. 25, 2002.

ACM set of points in $\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$ if and only if $\Delta H_{\mathbb{X}}$, a generalized first difference function, is the Hilbert function of some $\mathbb{N}^k$-graded artinian quotient. Our generalization relies on two main ingredients: (1) the existence of a regular sequence in $R/I_{\mathbb{X}}$ such that each element has a specific multi-degree, and (2) the techniques of [9] for lifting monomial ideals.

This characterization is not very satisfactory because it translates our original problem into the open problem of characterizing the Hilbert functions of $\mathbb{N}^k$-graded artinian quotients. However, we characterize these quotients in the special case $n_1 = \cdots = n_k = 1$, thereby giving a complete description of the Hilbert functions of ACM sets of points in $\mathbb{P}^1 \times \cdots \times \mathbb{P}^1$.

In the last two sections we specialize to ACM sets of points in $\mathbb{P}^1 \times \mathbb{P}^1$. It was first shown in [5] that the ACM sets of points are characterized by their Hilbert function. We give a new proof of this result, plus a new characterization that depends only upon numerical information describing the set $\mathbb{X}$. We then show that this numerical information also enables us to completely calculate the graded Betti numbers of the minimal free resolution of $I_{\mathbb{X}}$ (and thus, $H_{\mathbb{X}}$) provided $\mathbb{X}$ is ACM. This generalizes the fact that the Hilbert function and Betti numbers of a set of s points in $\mathbb{P}^1$ depend only upon s.

2. Preliminaries: Multi-graded rings, Hilbert functions, Points

Throughout this paper, $\mathbf{k}$ denotes an algebraically closed field of characteristic zero. In this section we provide the necessary facts and definitions about multi-graded rings, Hilbert functions, and sets of points in multi-projective spaces. See [10, 13] for more on these topics.

Let $\mathbb{N} := \{0, 1, 2, \ldots\}$. For an integer $n \in \mathbb{N}$, we set $[n] := \{1, \ldots, n\}$. We denote $(i_1, \ldots, i_k) \in \mathbb{N}^k$ by $\underline{i}$. We set $|\underline{i}| := \sum_h i_h$. If $\underline{i}, \underline{j} \in \mathbb{N}^k$, then $\underline{i} + \underline{j} := (i_1 + j_1, \ldots, i_k + j_k)$. We write $\underline{i} \geq \underline{j}$ if $i_h \geq j_h$ for every $h = 1, \ldots, k$. The set $\mathbb{N}^k$ is a semi-group generated by $\{e_1, \ldots, e_k\}$ where $e_i := (0, \ldots, 1, \ldots, 0)$ is the i^{th} standard basis vector of $\mathbb{N}^k$. If $c \in \mathbb{N}$, then $ce_i := (0, \ldots, c, \ldots, 0)$ with c in the i^{th} position.

Set $R = \mathbf{k}[x_{1,0}, \ldots, x_{1,n_1}, x_{2,0}, \ldots, x_{2,n_2}, \ldots, x_{k,0}, \ldots, x_{k,n_k}]$, and induce an $\mathbb{N}^k$-grading on R by setting $\deg x_{i,j} = e_i$. An element $x \in R$ is said to be $\mathbb{N}^k$-*homogeneous* (or simply *homogeneous* if the grading is clear) if $x \in R_{\underline{i}}$ for some $\underline{i} \in \mathbb{N}^k$. If x is homogeneous, then $\deg x := \underline{i}$. An ideal $I = (F_1, \ldots, F_r) \subseteq R$ is an $\mathbb{N}^k$-*homogeneous* (or simply, *homogeneous*) *ideal* if each F_j is $\mathbb{N}^k$-homogeneous.

For every $\underline{i} \in \mathbb{N}^k$, the set $R_{\underline{i}}$ is a finite dimensional vector space over $\mathbf{k}$. Since a basis for $R_{\underline{i}}$ is the set of all monomials of degree $\underline{i}$, $\dim_{\mathbf{k}} R_{\underline{i}} = \binom{n_1+i_1}{i_1}\binom{n_2+i_2}{i_2} \cdots \binom{n_k+i_k}{i_k}$.

If $I \subseteq R$ is a homogeneous ideal, then $S = R/I$ inherits an $\mathbb{N}^k$-graded ring structure if we define $S_{\underline{i}} = (R/I)_{\underline{i}} := R_{\underline{i}}/I_{\underline{i}}$. The numerical function $H_S : \mathbb{N}^k \rightarrow \mathbb{N}$ defined by $H_S(\underline{i}) := \dim_{\mathbf{k}}(R/I)_{\underline{i}} = \dim_{\mathbf{k}} R_{\underline{i}} - \dim_{\mathbf{k}} I_{\underline{i}}$ is the *Hilbert function of* S. If $H : \mathbb{N}^k \rightarrow \mathbb{N}$ is a numerical function, then the *first difference function of* H, denoted ΔH, is defined by

$$\Delta H(\underline{i}) := \sum_{\underline{0} \leq \underline{l} = (l_1, \ldots, l_k) \leq (1, \ldots, 1)} (-1)^{|\underline{l}|} H(i_1 - l_1, \ldots, i_k - l_k),$$

where $H(\underline{j}) = 0$ if $\underline{j} \not\geq \underline{0}$. If $k = 1$, then our definition agrees with the classical definition.

The $\mathbb{N}^k$-graded polynomial ring R is the coordinate ring of $\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$. Let

$$P = [a_{1,0} : \ldots : a_{1,n_1}] \times \cdots \times [a_{k,0} : \ldots : a_{k,n_k}] \in \mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$$

be a point in this space. The ideal of R associated to the point P is the prime ideal $I_P = (L_{1,1}, \ldots, L_{1,n_1}, \ldots, L_{k,1}, \ldots, L_{k,n_k})$ where $\deg L_{i,j} = e_i$ for $j = 1, \ldots, n_i$. If $P_1, \ldots, P_s$ are s distinct points and $\mathbb{X} = \{P_1, \ldots, P_s\} \subseteq \mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$, then $I_{\mathbb{X}} = I_{P_1} \cap \cdots \cap I_{P_s}$ where I_{P_i} is the ideal associated to the point P_i. The ring $R/I_{\mathbb{X}}$ then has the following property:

Lemma 2.1 ([12, Lemma 3.3])**.** *Let $\mathbb{X}$ be a finite set of points in $\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$. For each integer $i \in [k]$ there exists a form $L_i \in R_{e_i}$ such that $\overline{L}_i$ is a non-zero divisor in $R/I_{\mathbb{X}}$.*

We write $H_{\mathbb{X}}$ to denote the Hilbert function $H_{R/I_{\mathbb{X}}}$, and we say $H_{\mathbb{X}}$ is the *Hilbert function* of $\mathbb{X}$. Let $\pi_i : \mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k} \to \mathbb{P}^{n_i}$ denote the i^{th} projection morphism. Then $t_i := |\pi_i(\mathbb{X})|$ is the number of distinct i^{th}-coordinates in $\mathbb{X}$. With this notation we have

Proposition 2.2. *Let $\mathbb{X}$ be a finite set of points in $\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$ with Hilbert function $H_{\mathbb{X}}$.*

(*i*) [13, Proposition 4.2] *If $(j_1, \ldots, j_i, \ldots, j_k) \in \mathbb{N}^k$ and if $j_i \geq t_i - 1$, then*

$$H_{\mathbb{X}}(j_1, \ldots, j_i, \ldots, j_k) = H_{\mathbb{X}}(j_1, \ldots, t_i - 1, \ldots, j_k).$$

(*ii*) [13, Corollary 4.8] *$H_{\mathbb{X}}(j_1, \ldots, j_k) = s$ for all $(j_1, \ldots, j_k) \geq (t_1 - 1, t_2 - 1, \ldots, t_k - 1)$.*

Remark 2.3. Fix an integer $i \in [k]$, and fix $k-1$ integers in $\mathbb{N}$, say $j_1, \ldots, j_{i-1}, j_{i+1}, \ldots, j_k$. Set $\underline{j}_l := (j_1, \ldots, j_{i-1}, l, j_{i+1}, \ldots, j_k)$ for each integer $l \in \mathbb{N}$. Then Proposition 2.2 (i) can be interpreted as saying the sequence $\left\{H_{\mathbb{X}}(\underline{j}_l)\right\}$ becomes constant. In fact, $H_{\mathbb{X}}(\underline{j}_l) = H_{\mathbb{X}}(\underline{j}_{t_i-1})$ for all $l \geq t_i - 1$.

Proposition 2.4 ([13, Proposition 3.2])**.** *Let $\mathbb{X}$ be a finite set of points in $\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$ with Hilbert function $H_{\mathbb{X}}$. Fix an integer $i \in [k]$. Then the sequence $H = \{h_j\}$, where $h_j := H_{\mathbb{X}}(je_i)$, is the Hilbert function of $\pi_i(\mathbb{X}) \subseteq \mathbb{P}^{n_i}$.*

We end this section with some comments on the depth and Krull dimension of $R/I_{\mathbb{X}}$. Let $\mathbf{m} := \bigoplus_{0 \neq \underline{j} \in \mathbb{N}^k} R_{\underline{j}} = (x_{1,0}, \ldots, x_{k,n_k})$ be the maximal ideal of R. If $I \subseteq R$ is an $\mathbb{N}^k$-homogeneous ideal, then recall that we say a sequence $F_1, \ldots, F_r$ of elements is a *regular sequence modulo I* if and only if (*i*) $(I, F_1, \ldots, F_r) \subseteq \mathbf{m}$, (*ii*) $\overline{F}_1$ is not a zero divisor in R/I, and (*iii*) $\overline{F}_i$ is not a zero divisor in $R/(I, F_1, \ldots, F_{i-1})$ for $1 < i \leq r$. The *depth of R/I*, written $\operatorname{depth} R/I$, is the length of the maximal regular sequence modulo I.

Because each prime ideal I_{P_i} has height $\sum_{i=1}^{k} n_i$, it follows that $\text{K-dim}\, R/I_{\mathbb{X}} = k$, the number of projective spaces. This result, coupled with Lemma 2.1, implies $1 \leq \operatorname{depth} R/I_{\mathbb{X}} \leq k$. Thus, every set of points in $\mathbb{P}^n$ has $\operatorname{depth} R/I_{\mathbb{X}} = 1$. If $k \geq 2$, the value for $\operatorname{depth} R/I_{\mathbb{X}}$ is not immediately clear. In fact, for each integer $l \in [k]$ we can construct a set of points in $\mathbb{X}$ such that $\operatorname{depth} R/I_{\mathbb{X}} = l$. We begin with a lemma.

Lemma 2.5. *Fix a positive integer k. Denote by X_1 and X_2 the two points*

$$X_1 := [1:0] \times [1:0] \times \cdots \times [1:0] \text{ and } X_2 := [0:1] \times [0:1] \times \cdots \times [0:1]$$

in $\mathbb{P}^1 \times \cdots \times \mathbb{P}^1$ (k times). If $\mathbb{X} := \{X_1, X_2\}$, then $\operatorname{depth} R/I_{\mathbb{X}} = 1$.

Proof. The defining ideal of $\mathbb{X}$ is $I_{\mathbb{X}} = I_{X_1} \cap I_{X_2} = (\{x_{a,0}x_{b,1} \mid 1 \leq a \leq k, 1 \leq b \leq k\})$ in the $\mathbb{N}^k$-graded ring $R = \mathbf{k}[x_{1,0}, x_{1,1}, x_{2,0}, x_{2,1}, \ldots, x_{k,0}, x_{k,1}]$. Since $x_{1,0} + x_{1,1}$ does not vanish at either point, it suffices to show that every non-zero element of $R/(I_{\mathbb{X}}, x_{1,0} + x_{1,1})$ is a zero divisor.

So, set $J = (I_{\mathbb{X}}, x_{1,0} + x_{1,1})$ and suppose that $0 \neq \overline{F} \in R/J$. Without loss of generality, we can take F to be $\mathbb{N}^k$-homogeneous. We write F as $F = F_0 + F_1 x_{1,0} + F_2 x_{1,0}^2 + \cdots$ where $F_i \in$

$\mathbf{k}[x_{1,1},x_{2,0},\ldots,x_{k,1}]$. Since $x_{1,0}x_{1,1}\in I_{\mathbb{X}}$, it follows that $x_{1,0}^2 = x_{1,0}(x_{1,0}+x_{1,1}) - x_{1,0}x_{1,1}\in J$. Hence, we can assume that $F = F_0 + F_1x_{1,0}$. The element $x_{1,0}\notin J$. Although $x_{1,0}, F\notin J$, we claim that $Fx_{1,0}\in J$. Indeed, for each integer $1\leq b\leq k$, $x_{1,0}x_{b,1}\in I_{\mathbb{X}}\subseteq J$. Furthermore, for $1\leq a\leq k$, the element $x_{1,0}x_{a,0} = x_{a,0}\left(x_{1,0}+x_{1,1}\right) - x_{a,0}x_{1,1}\in J$. Hence, each term of $F_0x_{1,0}$ is in J, so $F_0x_{1,0}\in J$. But then $Fx_{1,0} = F_0x_{1,0} + F_1x_{1,0}^2\in J$. So, every $0\neq \overline{F}\in R/J$ is a zero divisor because it is annihilated by $\overline{x}_{1,0}$. □

Proposition 2.6. *Fix a positive integer k, and let $n_1,\ldots,n_k$ be k positive integers. Then, for every integer $l\in[k]$, there exists a set of points $\mathbb{X}$ in $\mathbb{P}^{n_1}\times\cdots\times\mathbb{P}^{n_k}$ with* $\operatorname{depth} R/I_{\mathbb{X}} = l$.

Proof. For every $l\in[k]$ we construct a set with the desired depth. Set $P_i := [1:0:\cdots:0]\in\mathbb{P}^{n_i}$ for $1\leq i\leq k$ and $Q_i := [0:1:0\cdots:0]\in\mathbb{P}^{n_i}$ for $1\leq i\leq k$. Fix an $l\in[k]$ and let X_1 and X_2 be the following two points of $\mathbb{P}^{n_1}\times\cdots\times\mathbb{P}^{n_k}$:

$$X_1 := P_1\times P_2\times\cdots\times P_k \text{ and } X_2 := P_1\times P_2\times\cdots\times P_{l-1}\times Q_l\times\ldots\times Q_k.$$

If we set $\mathbb{X}_l := \{X_1,X_2\}$, we claim that $\operatorname{depth} R/I_{\mathbb{X}_l} = l$. The defining ideal of $\mathbb{X}_l$ is

$$I_{\mathbb{X}_l} \;=\; \left(\begin{array}{l} x_{1,1},\ldots,x_{1,n_1},\ldots,x_{l-1,1},\ldots,x_{l-1,n_{l-1}},\\ x_{l,2},\ldots,x_{l,n_l},\ldots,x_{k,2},\ldots,x_{k,n_k},\\ \{x_{a,0}x_{b,1}\mid l\leq a\leq k,\ l\leq b\leq k\}\end{array}\right).$$

It follows that $R/I_{\mathbb{X}_l}\cong S/J$ where

$$S/J = \frac{\mathbf{k}[x_{1,0},x_{2,0},x_{3,0},\ldots,x_{l-1,0},x_{l,0},x_{l,1},x_{l+1,0},x_{l+1,1},\ldots,x_{k,0},x_{k,1}]}{(\{x_{a,0}x_{b,1}\mid l\leq a\leq k, l\leq b\leq k\})}.$$

The indeterminates $x_{1,0},x_{2,0},\ldots,x_{l-1,0}$ give rise to a regular sequence in S/J. Thus, $\operatorname{depth} R/I_{\mathbb{X}_l} = \operatorname{depth} S/J\geq l-1$. Set $K = (J,x_{1,0},\ldots,x_{l-1,0})$. Then

$$S/K\cong\frac{\mathbf{k}[x_{l,0},x_{l,1},x_{l+1,0},x_{l+1,1},\ldots,x_{k,0},x_{k,1}]}{(\{x_{a,0}x_{b,1}\mid l\leq a\leq k, l\leq b\leq k\})}.$$

The ring S/K is then isomorphic to the $\mathbb{N}^{k-l+1}$-graded coordinate ring of the two points $\{[1:0]\times[1:0]\times\cdots\times[1:0],[0:1]\times[0:1]\times\cdots\times[0:1]\}$ in $\mathbb{P}^1\times\cdots\times\mathbb{P}^1$ $((k-l+1)$ times). From Lemma 2.5 we have $\operatorname{depth} S/K = 1$, and hence, $\operatorname{depth} R/I_{\mathbb{X}_l} = l-1+1 = l$. □

3. The Hilbert functions of ACM Sets of Points

For an arbitrary set of points $\mathbb{X}\subseteq\mathbb{P}^{n_1}\times\cdots\times\mathbb{P}^{n_k}$, $\operatorname{depth} R/I_{\mathbb{X}}\leq \text{K-dim}\, R/I_{\mathbb{X}}$. If the equality holds, the coordinate ring is Cohen-Macaulay (CM), and the set of points are said to be arithmetically Cohen-Macaulay (ACM). We now investigate the Hilbert functions of those sets of points in $\mathbb{P}^{n_1}\times\cdots\times\mathbb{P}^{n_k}$ that are also ACM. Under this extra hypothesis we can generalize the characterization of the Hilbert functions of sets of points in $\mathbb{P}^n$ as found in [3].

We begin with a preparatory lemma.

Lemma 3.1. *Let $\mathbb{X}\subseteq\mathbb{P}^{n_1}\times\cdots\times\mathbb{P}^{n_k}$ be a finite set of points, and suppose $L_1,\ldots,L_t$, with $t\leq k$ and $\deg L_i = e_i$, give rise to a regular sequence in $R/I_{\mathbb{X}}$. Then there exists a positive integer l such that $(x_{1,0},\ldots,x_{1,n_1},\ldots,x_{t,0},\ldots,x_{t,n_t})^l\subseteq(I_{\mathbb{X}},L_1,\ldots,L_t)$.*

Proof. Set $J_i := (I_{\mathbb{X}},L_1,\ldots,L_i)$ for $i = 1,\ldots,t$. Since $L_1,\ldots,L_t$ form a regular sequence on $R/I_{\mathbb{X}}$, for each $i = 1,\ldots,t$ we have a short exact sequence with degree $(0,\ldots,0)$ maps:

$$0\rightarrow\left(R/J_{i-1}\right)(-e_i)\overset{\times\overline{L}_i}{\rightarrow} R/J_{i-1}\rightarrow R/J_i\rightarrow 0$$

where we set $J_0 := I_{\mathbb{X}}$. From the exact sequences we derive the following formula:

$$\dim_{\mathbf{k}}(R/J_t)_{\underline{i}} = \sum_{\underline{0} \leq (j_1,\ldots,j_t) \leq \underline{1}} (-1)^{(j_1+\cdots+j_t)} \dim_{\mathbf{k}}(R/I_{\mathbb{X}})_{i_1-j_1,\ldots,i_t-j_t,i_{t+1},\ldots,i_k}$$

where we set $\dim_{\mathbf{k}}(R/I_{\mathbb{X}})_{\underline{j}} = 0$ if $\underline{j} \not\geq \underline{0}$.

For each integer $j = 1,\ldots,t$, set $t_j := |\pi_j(\mathbb{X})|$. By Proposition 2.2, if $i_j \geq t_j$, then $\dim_{\mathbf{k}}(R/I_{\mathbb{X}})_{i_j e_j} = \dim_{\mathbf{k}}(R/I_{\mathbb{X}})_{(i_j-1)e_j}$. This fact, coupled with above above formula, implies that $\dim_{\mathbf{k}}(R/J_t)_{t_j e_j} = \dim_{\mathbf{k}}(R/I_{\mathbb{X}})_{t_j e_j} - \dim_{\mathbf{k}}(R/I_{\mathbb{X}})_{(t_j-1)e_j} = 0$. Thus $R_{t_j e_j} = (J_t)_{t_j e_j}$, or equivalently, $(x_{j,0},\ldots,x_{j,n_j})^{t_j} \subseteq (I_{\mathbb{X}}, L_1,\ldots,L_t)$. Since this is true for each integer $1 \leq j \leq t$, there exists an integer $l \gg 0$ such that $(x_{1,0},\ldots,x_{1,n_1},\ldots,x_{t,0},\ldots,x_{t,n_t})^l \subseteq J_t$, as desired. ☐

Proposition 3.2. *Suppose that $\mathbb{X}$ is an ACM set of points in $\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$. Then there exist elements $\overline{L}_1,\ldots,\overline{L}_k$ in $R/I_{\mathbb{X}}$ such that $L_1,\ldots,L_k$ give rise to a regular sequence in $R/I_{\mathbb{X}}$ and* $\deg L_i = e_i$.

Proof. The existence of a regular sequence of length k follows from the definition of a CM ring. The non-trivial part of this statement is the existence of a regular sequence whose elements have specific multi-degrees.

By Lemma 2.1 there exists a form $L_1 \in R_{e_1}$ such that $\overline{L}_1$ is a non-zero divisor of $R/I_{\mathbb{X}}$. To complete the proof it is enough to show for each $t = 2,...,k$ there exists an element $L_t \in R_{e_t}$ such that $\overline{L}_t$ is a non-zero divisor of the ring $R/(I_{\mathbb{X}}, L_1,\ldots,L_{t-1})$.

Set $J := (I_{\mathbb{X}}, L_1,\ldots,L_{t-1})$ and let $J = Q_1 \cap \cdots \cap Q_r$ be the ideal's primary decomposition. For each i set $\wp_i := \sqrt{Q_i}$. Since $\bigcup_{i=1}^r \overline{\wp}_i$ is the set of zero divisors of R/J, we want to show that $\bigcup_{i=1}^r (\wp_i)_{e_t} \subsetneq R_{e_t}$. Since R_{e_t} is a vector space over an infinite field, R_{e_t} cannot be expressed as the finite union of proper subvector spaces. Thus, it suffices to show $(\wp_i)_{e_t} \subsetneq R_{e_t}$ for each i.

So, suppose there is $i \in [r]$ such that $(\wp_i)_{e_t} = R_{e_t}$, or equivalently, $(x_{t,0},\ldots,x_{t,n_t}) \subseteq \wp_i$. By Lemma 3.1 there exists $l \in \mathbb{N}^+$ such that $(x_{1,0},\ldots,x_{1,n_1},\ldots,x_{t-1,0},\ldots,x_{t-1,n_{t-1}})^l \subseteq J \subseteq Q_i$. It follows that $(x_{1,0},\ldots,x_{1,n_1},\ldots,x_{t,0},\ldots,x_{t,n_t}) \subseteq \wp_i$.

Because the prime ideal $\wp_i$ also contains $I_{\mathbb{X}} = I_{P_1} \cap \cdots \cap I_{P_s}$, where I_{P_j} is the prime ideal associated to $P_j \in \mathbb{X}$, we can assume, after relabeling, $I_{P_1} \subseteq \wp_i$. Let $\wp := I_{P_1} + (x_{1,0},\ldots,x_{t,n_t})$. Since $I_{P_1} = (L_{1,1},\ldots,L_{1,n_1},\ldots,L_{k,1},\ldots,L_{k,n_k})$ where $\deg L_{m,n} = e_m$

$$\wp = (x_{1,0},\ldots,x_{t,n_t}, L_{t+1,1},\ldots,L_{t+1,n_{t+1}},\ldots,L_{k,1},\ldots,L_{k,n_k}) \subseteq \wp_i.$$

Thus $\mathrm{ht}_R(\wp_i) \geq \mathrm{ht}_R(\wp) = \left(\sum_{i=1}^k n_i\right) + t$, where $\mathrm{ht}_R(I)$ denotes the height of I.

From the identity $\mathrm{ht}_R(J) = \text{K-}\dim R - \text{K-}\dim R/J$ we calculate the height of J:

$$\mathrm{ht}_R(J) = \left(\sum_{i=1}^k n_i + 1\right) - (k-(t-1)) = \left(\sum_{i=1}^k n_i\right) + (t-1).$$

Since $\mathbb{X}$ is ACM, R/J is CM, and hence the ideal J is height unmixed, i.e., all the associated primes of J have height equal to $\mathrm{ht}_R(J)$. But $\wp_i$ is an associated prime of J with $\mathrm{ht}_R(\wp_i) > \mathrm{ht}_R(J)$. This contradiction implies our assumption $(\wp_i)_{e_t} = R_{e_t}$ cannot be true. ☐

Remark 3.3. If $S = \mathbf{k}[x_0,\ldots,x_n]$ is an $\mathbb{N}^1$-graded ring with $I \subseteq S$ such that S/I is CM, then a maximal regular sequence can be chosen so that each element is homogeneous [1, Proposition 1.5.11]. However, as stated in [11] (but no example is given), it is not always possible to pick a regular sequence that respects the multi-grading. For example, let $S = \mathbf{k}[x,y]$ with $\deg x = (1,0)$

and $\deg y = (0,1)$ and $I = (xy)$. Then S/I is CM, but all homogeneous elements of S/I, which have the form $\overline{cx}^a$ or $\overline{cy}^b$ with $c \in \mathbf{k}$, are zero divisors. Note that $\overline{x}+\overline{y}$ is a non-zero divisor, but not homogeneous. The fact that a homogeneous regular sequence can be found in a multi-graded ring is thus a very special situation.

We extend the notion of a graded artinian quotient in the natural way.

Definition 3.4. A homogeneous ideal I in the $\mathbb{N}^k$-graded ring R is an *artinian ideal* if any of the following equivalent statements hold:

(*i*) $\text{K-}\dim R/I = 0$.
(*ii*) $\sqrt{I} = \mathbf{m} = (x_{1,0},\ldots,x_{1,n_1},\ldots,x_{k,0},\ldots,x_{k,n_k})$.
(*iii*) For each integer $i \in [k]$, $H_{R/I}(le_i) = 0$ for all $l \gg 0$.

A ring $S = R/I$ is an $\mathbb{N}^k$-*graded artinian quotient* if the homogeneous ideal I is an artinian.

Corollary 3.5. *Let $\mathbb{X}$ be an ACM set of points in $\mathbb{P}^{n_1}\times\cdots\times\mathbb{P}^{n_k}$ with Hilbert function $H_{\mathbb{X}}$. Then*

$$\Delta H_{\mathbb{X}}(i_1,\ldots,i_k) := \sum_{\underline{0}\leq \underline{l}=(l_1,\ldots,l_k)\leq(1,\ldots,1)} (-1)^{|\underline{l}|} H_{\mathbb{X}}(i_1-l_1,\ldots,i_k-l_k),$$

where $H_{\mathbb{X}}(\underline{i}) = 0$ if $\underline{i} \not\geq \underline{0}$, is the Hilbert function of some $\mathbb{N}^k$-graded artinian quotient of the ring $\mathbf{k}[x_{1,1},\ldots,x_{1,n_1},\ldots,x_{k,1},\ldots,x_{k,n_k}]$.

Proof. By Proposition 3.2 there exists k forms $L_1,\ldots,L_k$ that give rise to a regular sequence in $R/I_{\mathbb{X}}$ with $\deg L_i = e_i$. After making a linear change of variables in the $x_{1,i}$'s, a linear change of variables in the $x_{2,i}$'s, etc., we can assume that $L_i = x_{i,0}$.

The ideal $(I_{\mathbb{X}},x_{1,0},\ldots,x_{k,0})/(x_{1,0},\ldots,x_{k,0})$ is isomorphic to an ideal J of the ring $S = \mathbf{k}[x_{1,1},\ldots,x_{1,n_1},\ldots,x_{k,1},\ldots,x_{k,n_k}]$. Set $A := S/J$, and so

$$A \cong \frac{R/(x_{1,0},\ldots,x_{k,0})}{(I_{\mathbb{X}},x_{1,0},\ldots,x_{k,0})/(x_{1,0},\ldots,x_{k,0})} \cong \frac{R}{(I_{\mathbb{X}},x_{1,0},\ldots,x_{k,0})}.$$

The ring A is artinian because there exists $l \gg 0$ by Lemma 3.1 such that $\mathbf{m}^l \subseteq (I_{\mathbb{X}},x_{1,0},\ldots,x_{k,0})$.

It therefore remains to compute the Hilbert function of A. Set $J_i = (I_{\mathbb{X}},x_{1,0},\ldots,x_{i,0})$ for $i = 1,\ldots,k$. For each $i = 1,\ldots,k$ we have a short exact sequence with degree $(0,\ldots,0)$ maps:

$$0 \longrightarrow (R/J_{i-1})(-e_i) \overset{\times\overline{x}_{i,0}}{\longrightarrow} R/J_{i-1} \longrightarrow R/J_i \longrightarrow 0$$

where $J_0 := I_{\mathbb{X}}$. From the k short exact sequences we have that

$$H_{R/J_k}(\underline{i}) = \Delta H_{\mathbb{X}}(\underline{i}) := \sum_{\underline{0}\leq \underline{l}=(l_1,\ldots,l_k)\leq(1,\ldots,1)} (-1)^{|\underline{l}|} H_{\mathbb{X}}(i_1-l_1,\ldots,i_k-l_k)$$

where $H_{\mathbb{X}}(\underline{i}) = 0$ if $\underline{i} \not\geq \underline{0}$. This completes the proof since $A \cong R/J_k$. □

The remainder of this section is devoted to showing the necessary condition in Corollary 3.5 is also sufficient. To demonstrate this converse, we describe how to *lift* an ideal.

Definition 3.6. Let $R = \mathbf{k}[x_{1,0},\ldots,x_{1,n_1},\ldots,x_{k,0},\ldots,x_{k,n_k}]$ and let $S = \mathbf{k}[x_{1,1},\ldots,x_{1,n_1},\ldots,x_{k,1},\ldots,x_{k,n_k}]$ be $\mathbb{N}^k$-graded rings. Let $I \subseteq R$ and $J \subseteq S$ be $\mathbb{N}^k$-homogeneous ideals. Then we say I is a *lifting of J to R* if

(*i*) I is radical in R

(*ii*) $\dfrac{(I, x_{1,0}, \ldots, x_{k,0})}{(x_{1,0}, \ldots, x_{k,0})} \cong J$

(*iii*) $x_{1,0}, \ldots, x_{k,0}$ give rise to a regular sequence in R/I.

Our plan is to lift a monomial ideal of S to an $\mathbb{N}^k$-homogeneous ideal I of R, using the techniques and results of [9], so such that I is the ideal of a reduced set of points in $\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$. We make a brief digression to introduce the relevant content of [9].

Suppose that S and R are as in Definition 3.6, but for the moment, we only assume that they are $\mathbb{N}^1$-graded. For each indeterminate $x_{i,j}$ with $1 \leq i \leq k$ and $1 \leq j \leq n_i$, choose infinitely many linear forms $L_{i,j,l} \in \mathbf{k}[x_{i,j}, x_{1,0}, x_{2,0}, \ldots, x_{k,0}]$ with $l \in \mathbb{N}^+$. We only assume that the coefficient of $x_{i,j}$ in $L_{i,j,l}$ is not zero. The infinite matrix A, where

$$A := \begin{bmatrix} L_{1,1,1} & L_{1,1,2} & L_{1,1,3} & \cdots \\ \vdots & \vdots & \vdots & \\ L_{1,n_1,1} & L_{1,n_1,2} & L_{1,n_1,3} & \cdots \\ \vdots & \vdots & \vdots & \\ L_{k,1,1} & L_{k,1,2} & L_{k,1,3} & \cdots \\ \vdots & \vdots & \vdots & \\ L_{k,n_k,1} & L_{k,n_k1,2} & L_{k,n_k,3} & \cdots \end{bmatrix}$$

is called a *lifting matrix.* By using the lifting matrix, we associate to each monomial $m = x_{1,1}^{a_{1,1}} \cdots x_{1,n_1}^{a_{1,n_1}} \cdots x_{k,1}^{a_{k,1}} \cdots x_{k,n_k}^{a_{k,n_k}}$ of the ring S the element

$$\overline{m} = \left[\prod_{i=1}^{n_1}\left(\prod_{j=1}^{a_{1,i}} L_{1,i,j}\right)\right] \cdots \left[\prod_{i=1}^{n_k}\left(\prod_{j=1}^{a_{k,i}} L_{k,i,j}\right)\right] \in R.$$

Depending upon our choice of $L_{i,j,l}$'s, $\overline{m}$ may or may not be $\mathbb{N}^k$-homogeneous. However, $\overline{m}$ is homogeneous. If $J = (m_1, \ldots, m_r)$ is a monomial ideal of S, then we use I to denote the ideal $(\overline{m}_1, \ldots, \overline{m}_r) \subseteq R$. The following properties, among others, relate R/I and S/J.

Proposition 3.7 ([9, Corollary 2.10]). *Let $J \subseteq S$ be a monomial ideal, and let I be the ideal constructed from J using any lifting matrix. Then*

(*i*) *S/J is CM if and only if R/I is CM.*

(*ii*) $\dfrac{(I, x_{1,0}, \ldots, x_{k,0})}{(x_{1,0}, \ldots, x_{k,0})} \cong J$

(*iii*) $x_{1,0}, \ldots, x_{k,0}$ *give rise to a regular sequence in R/I.*

We now consider the lifting of a monomial ideal using the lifting matrix $\mathcal{A} := [L_{i,j,l}]$ where

$$L_{i,j,l} = x_{i,j} - (l-1)x_{i,0} \text{ for } 1 \leq i \leq n,\ 1 \leq j \leq n_i, \text{ and } l \in \mathbb{N}^+.$$

The lifting matrix $\mathcal{A}$ associates to every monomial $m = x_{1,1}^{a_{1,1}} \cdots x_{1,n_1}^{a_{1,n_1}} \cdots x_{k,1}^{a_{k,1}} \cdots x_{k,n_k}^{a_{k,n_k}}$ of S the following $\mathbb{N}^k$-homogeneous form of R:

$$\overline{m} = \left[\prod_{i=1}^{n_1}\left(\prod_{j=1}^{a_{1,i}}(x_{1,i} - (j-1)x_{1,0})\right)\right] \cdots \left[\prod_{i=1}^{n_k}\left(\prod_{j=1}^{a_{k,i}}(x_{k,i} - (j-1)x_{k,0})\right)\right].$$

Thus, if I is constructed from a monomial ideal $J \subseteq S$ using $\mathcal{A}$, I is $\mathbb{N}^k$-homogeneous. In fact, using $\mathcal{A}$, the ideal I is a lifting of J to R. To prove this statement, we require

Lemma 3.8 ([9, Corollary 2.18])**.** *Let $J \subseteq S$ be a monomial ideal and let I be be constructed from J using any lifting matrix. If $J = Q_1 \cap \cdots \cap Q_r$ is the primary decomposition of J, then $I = \overline{Q}_1 \cap \cdots \cap \overline{Q}_r$ where $\overline{Q}_i$ is the ideal generated by the lifting of the generators of Q_i.*

Proposition 3.9. *Let $J \subseteq S$ be a monomial ideal and let I be the ideal constructed from J using the lifting matrix $\mathcal{A}$. Then I is a lifting of J to R.*

Proof. Since Proposition 3.7 is true for any lifting matrix, it suffices to show that I is radical. Let $J = Q_1 \cap \cdots \cap Q_r$ be the primary decomposition of J. Since J is a monomial ideal, then by Remark 2.19 of [9] we have that each Q_i is a complete intersection of the form

$$Q_i = \left(x_{1,i_{1,1}}^{a_{1,i_{1,1}}}, \ldots, x_{1,i_{1,p_1}}^{a_{1,i_{1,p_1}}}, \ldots x_{k,i_{k,1}}^{a_{k,i_{k,1}}}, \ldots, x_{k,i_{k,p_k}}^{a_{k,i_{k,p_k}}}\right)$$

with $a_{j,i_{j,l}} \geq 1$ for each variable that appears in Q_i. Using the lifting matrix $\mathcal{A}$ we then have

$$\overline{Q}_i = \left(\prod_{l=1}^{a_{1,i_{1,1}}} L_{1,i_{1,1},l}, \ldots, \prod_{l=1}^{a_{1,i_{1,p_1}}} L_{1,i_{1,p_1},l}, \ldots, \prod_{l=1}^{a_{k,i_{k,1}}} L_{k,i_{k,1},l}, \ldots, \prod_{l=1}^{a_{k,i_{k,p_k}}} L_{k,i_{k,p_k},l}\right)$$

where $L_{i,j,l} = x_{i,j} - (l-1)x_{i,0}$. But then $\overline{Q}_i$ is a reduced complete intersection. It then follows from Lemma 3.8 that I must be radical. $\square$

We now describe the zero set of the lifted ideal I. For each

$$(\underline{\alpha}_1, \ldots, \underline{\alpha}_k) := ((a_{1,1}, \ldots, a_{1,n_1}), \ldots, (a_{k,1}, \ldots, a_{k,n_k})) \in \mathbb{N}^{n_1} \times \cdots \times \mathbb{N}^{n_k},$$

set $X_1^{\underline{\alpha}_1} \cdots X_k^{\underline{\alpha}_k} := x_{1,1}^{a_{1,1}} \cdots x_{1,n_1}^{a_{1,n_1}} \cdots x_{k,1}^{a_{k,1}} \cdots x_{k,n_k}^{a_{k,n_k}}$. If P is the set of all monomials of S including the monomial 1, then there exists a bijection between P and $\mathbb{N}^{n_1} \times \cdots \times \mathbb{N}^{n_k}$ given by the map $X_1^{\underline{\alpha}_1} \cdots X_k^{\underline{\alpha}_k} \leftrightarrow (\underline{\alpha}_1, \ldots, \underline{\alpha}_k)$. To each tuple $(\underline{\alpha}_1, \ldots, \underline{\alpha}_k)$ we associate the point $\overline{(\underline{\alpha}_1, \ldots, \underline{\alpha}_k)} \in \mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$ where

$$\overline{(\underline{\alpha}_1, \ldots, \underline{\alpha}_k)} := [1 : a_{1,1} : a_{1,2} : \cdots : a_{1,n_1}] \times \cdots \times [1 : a_{k,1} : a_{k,2} : \cdots : a_{k,n_k}].$$

Note that if $m = X_1^{\underline{\alpha}_1} \cdots X_k^{\underline{\alpha}_k} \in P$ and if $\overline{m}$ is constructed from $X_1^{\underline{\alpha}_1} \cdots X_k^{\underline{\alpha}_k}$ using the lifting matrix $\mathcal{A}$, then $\overline{m}(\overline{(\underline{\alpha}_1, \ldots, \underline{\alpha}_k)}) \neq 0$. In fact, it follows from our construction that $\overline{m}(\overline{(\underline{\beta}_1, \ldots, \underline{\beta}_k)}) = 0$ if and only if some coordinate of $(\underline{\beta}_1, \ldots, \underline{\beta}_k)$ is strictly less than some coordinate of $(\underline{\alpha}_1, \ldots, \underline{\alpha}_k)$.

If J is a monomial ideal of S, then let N be the set of monomials in J. The elements of $M := P \backslash N$ are representatives for a $\mathbf{k}$-basis of the $\mathbb{N}^k$-graded ring S/J. Set

$$\overline{M} := \left\{ \overline{(\underline{\beta}_1, \ldots, \underline{\beta}_k)} \in \mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k} \;\middle|\; X_1^{\underline{\beta}_1} \cdots X_k^{\underline{\beta}_k} \in M \right\}.$$

Let $\mathbf{I}(\overline{M})$ denote the $\mathbb{N}^k$-homogeneous ideal associated to $\overline{M}$. If $m = X_1^{\underline{\alpha}_1} \cdots X_k^{\underline{\alpha}_k} \in J$ is a minimal generator, then for each $X_1^{\underline{\beta}_1} \cdots X_k^{\underline{\beta}_k} \in M$ there exists at least one coordinate of $(\underline{\beta}_1, \ldots, \underline{\beta}_k)$ that is strictly less then some coordinate of $(\underline{\alpha}_1, \ldots, \underline{\alpha}_k)$. So $\overline{m}(\overline{(\underline{\beta}_1, \ldots, \underline{\beta}_k)}) = 0$ for all $X_1^{\underline{\beta}_1} \cdots X_k^{\underline{\beta}_k} \in M$, and hence the lifted ideal $I \subseteq \mathbf{I}(\overline{M})$. On the other hand $\overline{M} = \mathbf{V}(I)$, the zero set of I, so by the $\mathbb{N}^k$-graded analog of the Nullstellensatz (cf. [13, Theorem 2.3]) we have $\mathbf{I}(\overline{M}) \subseteq \sqrt{I}$. Because I is radical by Proposition 3.9 we have just shown

Lemma 3.10. *Let $J \subseteq S$ be a monomial ideal and let I be the ideal constructed from J using the lifting matrix $\mathcal{A}$. Then, with the notation as above, $I = \mathbf{I}(\overline{M})$.*

We come to the main result of this section.

Theorem 3.11. *Let $H : \mathbb{N}^k \to \mathbb{N}$ be a numerical function. Then H is the Hilbert function of an ACM set of distinct points in $\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$ if and only if the first difference function*

$$\Delta H(i_1, \ldots, i_k) = \sum_{\underline{0} \leq \underline{l} = (l_1, \ldots, l_k) \leq (1, \ldots, 1)} (-1)^{|\underline{l}|} H(i_1 - l_1, \ldots, i_k - l_k),$$

where $H(\underline{i}) = 0$ if $\underline{i} \ngeq \underline{0}$, is the Hilbert function of some $\mathbb{N}^k$-graded artinian quotient of $S = \mathbf{k}[x_{1,1}, \ldots, x_{1,n_1}, \ldots, x_{k,1}, \ldots, x_{k,n_k}]$.

Proof. Because of Corollary 3.5, we only need to show one direction. So, if ΔH is the Hilbert function of some $\mathbb{N}^k$-graded artinian quotient of S, then there exists an $\mathbb{N}^k$-homogeneous ideal $J \subseteq S$ with $\Delta H(\underline{i}) = H_{S/J}(\underline{i})$ for all $\underline{i} \in \mathbb{N}^k$. By replacing J with its leading term ideal, we can assume that $J = (m_1, \ldots, m_r)$ is a monomial ideal of S.

Let $I \subseteq R$ be the ideal constructed from J using the lifting matrix $\mathcal{A}$. By Proposition 3.9, the ideal $J \cong (I, x_{1,0}, \ldots, x_{k,0})/(x_{1,0}, \ldots, x_{k,0})$ where $x_{1,0}, \ldots, x_{k,0}$ give rise to a regular sequence in R/I. Because $\deg x_{i,0} = e_i$, we have k short exact sequences with degree $(0, \ldots, 0)$ maps:

$$0 \longrightarrow (R/J_{i-1})\,(e_i) \xrightarrow{\times \overline{x}_{i,0}} R/J_{i-1} \longrightarrow R/J_i \longrightarrow 0$$

where $J_i := (I, x_{1,0}, \ldots, x_{i,0})$ for $i = 1, \ldots, k$ and $J_0 := I$. Furthermore,

$$S/J \cong \frac{R/(x_{1,0}, \ldots, x_{k,0})}{(I, x_{1,0}, \ldots, x_{k,0})/(x_{1,0}, \ldots, x_{k,0})} \cong R/(I, x_{1,0}, \ldots, x_{k,0}) = R/J_k.$$

Then, using the k short exact sequences to calculate the $H_{R/I}$, we find that $H = H_{R/I}$.

If N is the set of monomials in J, then $M = P \backslash N$ is a finite set of monomials because S/J is artinian. By Lemma 3.10, I is the ideal of the finite set of points $\overline{M} \subseteq \mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$. Finally, by Proposition 3.7 the set $\overline{M}$ is ACM because S/J is artinian, and hence, CM. ☐

Since characterizing the Hilbert functions of ACM sets of points in $\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$ is equivalent to characterizing the Hilbert functions of $\mathbb{N}^k$-graded artinian quotients of S, Theorem 3.11 translates one open problem into another open problem because we do not have a theorem like Macaulay's Theorem [7] for $\mathbb{N}^k$-graded rings if $k > 1$. However, as shown below, there is a Macaulay-type theorem for $\mathbb{N}^k$-graded quotients of $\mathbf{k}[x_{1,1}, x_{2,1}, \ldots, x_{k,1}]$. As a consequence, we can explicitly describe all the Hilbert functions of ACM sets of points in $\mathbb{P}^1 \times \cdots \times \mathbb{P}^1$ (k times) for any positive integer k.

So, suppose that $S = \mathbf{k}[x_1, \ldots, x_k]$ and $\deg x_i = e_i$, where e_i is the i^{th} standard basis vector of $\mathbb{N}^k$. We prove a stronger result by characterizing the Hilbert functions of all quotients of S, not only the artinian quotients.

Theorem 3.12. *Let $S = \mathbf{k}[x_1, \ldots, x_k]$ with $\deg x_i = e_i$, and let $H : \mathbb{N}^k \to \mathbb{N}$ be a numerical function. Then there exists a homogeneous ideal $I \subsetneq S$ with Hilbert function $H_{S/I} = H$ if and only if*

- (*i*) $H(0, \ldots, 0) = 1$,
- (*ii*) $H(\underline{i}) = 1$ *or* 0 *if* $\underline{i} > \underline{0}$, *and*
- (*iii*) *if* $H(\underline{i}) = 0$, *then* $H(\underline{j}) = 0$ *for all* $\underline{j} \geq \underline{i}$.

Proof. If $I \subsetneq S$ is an $\mathbb{N}^k$-homogeneous ideal such that $H_{S/I} = H$, then condition (*i*) is simply a consequence of the fact that $I \subsetneq S$. Statement (*ii*) follows from the inequality $0 \leq H_{S/I}(\underline{i}) \leq$

$\dim_{\mathbf{k}} S_{\underline{i}} = 1$. Finally, if $H_{S/I}(\underline{i}) = 0$, then $x_1^{i_1} \cdots x_k^{i_k} \in I$, or equivalently, $S_{\underline{i}} \subseteq I$ because $x_1^{i_1} \cdots x_k^{i_k}$ is a basis for $S_{\underline{i}}$. So, if $\underline{j} \geq \underline{i}$, then $S_{\underline{j}} \subseteq I$, and hence, $H_{S/I}(\underline{j}) = 0$, thus proving (iii).

Conversely, suppose that H is a numerical function satisfying $(i) - (iii)$. If $H(\underline{i}) = 1$ for all $\underline{i} \in \mathbb{N}^k$, then the ideal $I = (0) \subseteq S$ has the property that $H_{S/I} = H$.

So, suppose $H(\underline{i}) \neq 1$ for all $\underline{i}$. Set $\mathcal{I} := \left\{\underline{i} \in \mathbb{N}^k \mid H(\underline{i}) = 0\right\}$. Note that $\mathcal{I} \neq \mathbb{N}^k$ because $\underline{0} \notin \mathcal{I}$. Let I be the ideal $I := \langle \{x_1^{i_1} \cdots x_k^{i_k} \mid \underline{i} \in \mathcal{I}\}\rangle$ in S. We claim that $H_{S/I}(\underline{i}) = H(\underline{i})$ for all $\underline{i} \in \mathbb{N}^k$. It is immediate that $H_{S/I}(\underline{0}) = H(\underline{0}) = 1$. Moreover, if $H(\underline{i}) = 0$, then $H_{S/I}(\underline{i}) = 0$ because $x_1^{i_1} \cdots x_k^{i_k} \in I_{\underline{i}} \subseteq I$, i.e., $S_{\underline{i}} \subseteq I$.

So, we need to check that if $H(\underline{i}) = 1$, then $H_{S/I}(\underline{i}) = 1$. Suppose $H_{S/I}(\underline{i}) = 0$. This implies that $x_1^{i_1} \cdots x_k^{i_k} \in I$. But because $\underline{i} \notin \mathcal{I}$, there is a monomial $x_1^{j_1} \cdots x_k^{j_k} \in I$ with $\underline{j} \in \mathcal{I}$, such that $x_1^{j_1} \cdots x_k^{j_k}$ divides $x_1^{i_1} \cdots x_k^{i_k}$. But this is equivalent to the statement that $\underline{j} \leq \underline{i}$. But this contradicts hypothesis (iii). So $H_{S/I}(\underline{i}) = 1$. □

Using Theorem 3.12 and the definition of an $\mathbb{N}^k$-graded artinian quotient we then have:

Corollary 3.13. *Let $S = \mathbf{k}[x_1, \ldots, x_k]$ with $\deg x_i = e_i$, and let $H : \mathbb{N}^k \to \mathbb{N}$ be a numerical function. Then H is the Hilbert function of an $\mathbb{N}^k$-graded artinian quotient of S if and only if*

(i) $H(0, \ldots, 0) = 1$,
(ii) $H(\underline{i}) = 1$ *or* 0 *if* $\underline{i} > (0, \ldots, 0)$,
(iii) *if* $H(\underline{i}) = 0$, *then* $H(\underline{j}) = 0$ *for all* $\underline{j} \geq \underline{i}$, *and*
(iv) *for each* $i \in [k]$ *there exists an integer* t_i *such that* $H(t_i e_i) = 0$.

Corollary 3.14. *Let $H : \mathbb{N}^k \to \mathbb{N}$ be a numerical function. Then H is the Hilbert function of an ACM set of distinct points in $\mathbb{P}^1 \times \cdots \times \mathbb{P}^1$ (k times) if and only ΔH satisfies conditions $(i) - (iv)$ of Corollary 3.13.*

Remark 3.15. It follows from the previous corollaries that H is the Hilbert function of an ACM set of points in $\mathbb{P}^1 \times \mathbb{P}^1$ if and only if

(i) $\Delta H(i, j) = 1$ or 0,
(ii) if $\Delta H(i, j) = 0$, then $\Delta H(i', j') = 0$ for all $(i', j') \in \mathbb{N}^2$ with $(i', j') > (i, j)$, and
(iii) there exists integers t and r such that $\Delta H(t, 0) = 0$ and $\Delta H(0, r) = 0$.

Giuffrida, Maggioni, and Ragusa proved precisely this result in Theorems 4.1 and 4.2 of [5]. We investigate ACM sets of points in $\mathbb{P}^1 \times \mathbb{P}^1$ in further detail in the next two sections.

4. ACM Sets of Points in $\mathbb{P}^1 \times \mathbb{P}^1$

If $\mathbb{X}$ is an ACM set of points in $\mathbb{P}^1 \times \mathbb{P}^1$, then by Theorem 3.11 the function $\Delta H_{\mathbb{X}}$ is the Hilbert function of a bigraded artinian quotient of $\mathbf{k}[x_1, y_1]$. In [5] it was shown that the converse of this statement is also true, thereby classifying the ACM sets of points in $\mathbb{P}^1 \times \mathbb{P}^1$. In this section we revisit this result by giving a new proof of this characterization that depends only upon numerical information describing $\mathbb{X}$.

We begin with a brief digression to introduce some needed combinatorial results. Recall that a tuple $\lambda = (\lambda_1, \ldots, \lambda_r)$ of positive integers is a *partition* of an integer s, denoted $\lambda \vdash s$, if $\sum \lambda_j = s$ and $\lambda_i \geq \lambda_{i+1}$ for each i. If $\lambda \vdash s$, then the *conjugate* of λ is the tuple $\lambda^* = (\lambda_1^*, \ldots, \lambda_{\lambda_1}^*)$ where $\lambda_i^* := \#\{\lambda_j \in \lambda \mid \lambda_j \geq i\}$. Moreover, λ^* is also a partition of s.

To any partition $\lambda=(\lambda_1,\ldots,\lambda_r)\vdash s$ we can associate the following diagram: on an $r\times\lambda_1$ grid, place λ_1 points on the first line, λ_2 points on the second, and so on. The resulting diagram is called the *Ferrers diagram* of λ. For example, suppose $\lambda=(4,4,3,1,1)\vdash 13$. Then the Ferrers diagram is

$$\begin{array}{cccc}\bullet&\bullet&\bullet&\bullet\\ \bullet&\bullet&\bullet&\bullet\\ \bullet&\bullet&\bullet&\\ \bullet&&&\\ \bullet&&&\end{array}$$

The conjugate of λ can be read off the Ferrers diagram by counting the number of dots in each column as opposed to each row. In this example $\lambda^*=(5,3,3,2)\vdash 13$.

The following lemma, whose proof is a straightforward combinatorial exercise, describes some of the relations between a partition and its conjugate.

Lemma 4.1. *Let* $\alpha=(\alpha_1,\ldots,\alpha_n)\vdash s$, *and* $\beta=(\beta_1,\ldots,\beta_m)\vdash s$. *If* $\alpha^*=\beta$, *then*

(*i*) $\alpha_1=|\beta|$ *and* $\beta_1=|\alpha|$.
(*ii*) *if* $\alpha'=(\alpha_2,\ldots,\alpha_n)$ *and* $\beta'=(\beta_1-1,\ldots,\beta_{\alpha_2}-1)$, *then* $(\alpha')^*=\beta'$.

Let $\mathbb{X}$ denote a set of reduced points in $\mathbb{P}^1\times\mathbb{P}^1$, and associate to $\mathbb{X}$ two tuples $\alpha_{\mathbb{X}}$ and $\beta_{\mathbb{X}}$ as follows. Let $\pi_1(\mathbb{X})=\{P_1,\ldots,P_t\}$ be the t distinct first coordinates in $\mathbb{X}$. Then, for each $P_i\in\pi_1(\mathbb{X})$, let $\alpha_i:=|\pi_1^{-1}(P_i)|$, i.e., the number of points in $\mathbb{X}$ which have P_i as its first coordinate. After relabeling the α_i so that $\alpha_i\geq\alpha_{i+1}$ for $i=1,\ldots,t-1$, we set $\alpha_{\mathbb{X}}=(\alpha_1,\ldots,\alpha_t)$. Analogously, for each $Q_i\in\pi_2(\mathbb{X})=\{Q_1,\ldots,Q_r\}$, we let $\beta_i:=|\pi_2^{-1}(Q_i)|$. After relabeling so that $\beta_i\geq\beta_{i+1}$ for $i=1,\ldots,r-1$, we set $\beta_{\mathbb{X}}=(\beta_1,\ldots,\beta_r)$. So, by construction, $\alpha_{\mathbb{X}},\beta_{\mathbb{X}}\vdash s=|\mathbb{X}|$. Note that $|\alpha_{\mathbb{X}}|=|\pi_1(\mathbb{X})|$ and $|\beta_{\mathbb{X}}|=|\pi_2(\mathbb{X})|$.

We write the Hilbert function $H_{\mathbb{X}}$ as an infinite matrix (m_{ij}) where $m_{ij}:=H_{\mathbb{X}}(i,j)$. Proposition 2.4 gives

$$m_{i,0}=\begin{cases} i+1 & 0\leq i\leq t-1\\ t & i\geq t\end{cases}\quad\text{and } m_{0,j}=\begin{cases} i+1 & 0\leq i\leq r-1\\ r & i\geq r\end{cases}.$$

because $\pi_1(\mathbb{X})=\{P_1,\ldots,P_t\}\subseteq\mathbb{P}^1$ and $\pi_2(\mathbb{X})=\{Q_1,\ldots,Q_r\}\subseteq\mathbb{P}^1$. This fact, combined with Proposition 2.2, implies that $H_{\mathbb{X}}$ has the form

$$(1)\qquad H_{\mathbb{X}}=\begin{bmatrix} 1 & 2 & \cdots & r-1 & \mathbf{r} & r & \cdots\\ 2 & & & & \mathbf{m_{1,r-1}} & m_{1,r-1} & \cdots\\ \vdots & & * & & \vdots & \vdots & \\ t-1 & & & & \mathbf{m_{2,r-1}} & m_{2,r-1} & \cdots\\ \mathbf{t} & \mathbf{m_{t-1,1}} & \cdots & \mathbf{m_{t-1,r-2}} & \mathbf{s} & s & \cdots\\ t & m_{t-1,1} & \cdots & m_{t-1,r-2} & s & s & \\ \vdots & \vdots & & \vdots & \vdots & & \ddots\end{bmatrix}$$

where the values denoted by $(*)$ need to be calculated.

Set $B_C=(m_{t-1,0},\ldots,m_{t-1,r-1})$ and $B_R=(m_{0,r-1},\ldots,m_{t-1,r-1})$. From our description of $H_{\mathbb{X}}$, we see that if we know the values in the tuples B_C and B_R, we will know all but a finite number of values of $H_{\mathbb{X}}$. As shown below, the tuples B_C and B_R can be computed directly from the tuples $\alpha_{\mathbb{X}}$ and $\beta_{\mathbb{X}}$ defined above. If λ is a tuple, then we shall abuse notation and write $\lambda_i\in\lambda$ to mean that λ_i is a coordinate of λ.

Proposition 4.2 ([13, Proposition 5.11])**.** *Let $\mathbb{X} \subseteq \mathbb{P}^1 \times \mathbb{P}^1$ be a finite set of points in $\mathbb{P}^1 \times \mathbb{P}^1$.*

(*i*) *if $B_C = (m_{t-1,0}, \ldots, m_{t-1,r-1})$ where $m_{t-1,j} = H_{\mathbb{X}}(t-1,j)$ for $j = 0, \ldots, r-1$, then*
$$m_{t-1,j} = \#\{\alpha_i \in \alpha_{\mathbb{X}} \mid \alpha_i \geq 1\} + \#\{\alpha_i \in \alpha_{\mathbb{X}} \mid \alpha_i \geq 2\} + \cdots + \#\{\alpha_i \in \alpha_{\mathbb{X}} \mid \alpha_i \geq j+1\}.$$

(*ii*) *if $B_R = (m_{0,r-1}, \ldots, m_{t-1,r-1})$ where $m_{j,r-1} = H_{\mathbb{X}}(j,r-1)$ for $j = 0, \ldots, t-1$, then*
$$m_{j,r-1} = \#\{\beta_i \in \beta_{\mathbb{X}} \mid \beta_i \geq 1\} + \#\{\beta_i \in \beta_{\mathbb{X}} \mid \beta_i \geq 2\} + \cdots + \#\{\beta_i \in \beta_{\mathbb{X}} \mid \beta_i \geq j+1\}.$$

We can rewrite this result more compactly using the language of combinatorics introduced above. If $p = (p_1, \ldots, p_k)$, then we write Δp to denote the tuple $\Delta p := (p_1, p_2 - p_1, \ldots, p_k - p_{k-1})$.

Corollary 4.3 ([13, Corollary 5.13])**.** *Let $\mathbb{X}$ be a finite set of points in $\mathbb{P}^1 \times \mathbb{P}^1$. Then*

(*i*) $\Delta B_C = \alpha^*_{\mathbb{X}}$.
(*ii*) $\Delta B_R = \beta^*_{\mathbb{X}}$.

Remark 4.4. In [13] the tuple $B_{\mathbb{X}} = (B_C, B_R)$ was called the *border* of the Hilbert function.

Recall that $\Delta H_{\mathbb{X}}$, the *first difference function* of $H_{\mathbb{X}}$, is defined by
$$\Delta H_{\mathbb{X}}(i,j) := H_{\mathbb{X}}(i,j) - H_{\mathbb{X}}(i-1,j) - H_{\mathbb{X}}(i,j-1) + H_{\mathbb{X}}(i-1,j-1)$$
where $H_{\mathbb{X}}(i,j) = 0$ if $(i,j) \not\succeq (0,0)$. The entries of $\alpha^*_{\mathbb{X}}$ and $\beta^*_{\mathbb{X}}$ can then be read from $\Delta H_{\mathbb{X}}$.

Corollary 4.5. *Let $\mathbb{X} \subseteq \mathbb{P}^1 \times \mathbb{P}^1$ be a finite set of points, and set $c_{i,j} := \Delta H_{\mathbb{X}}(i,j)$. Then*

(*i*) *for every $0 \leq j \leq r-1 = |\pi_2(\mathbb{X})| - 1$*
$$\alpha^*_{j+1} = \sum_{h \leq |\pi_1(\mathbb{X})|-1} c_{h,j}$$
where α^*_{j+1} is the $(j+1)^{th}$ entry of $\alpha^*_{\mathbb{X}}$, the conjugate of $\alpha_{\mathbb{X}}$.

(*ii*) *for every $0 \leq i \leq t-1 = |\pi_1(\mathbb{X})| - 1$*
$$\beta^*_{i+1} = \sum_{h \leq |\pi_2(\mathbb{X})|-1} c_{i,h}.$$
where β^*_{i+1} is the $(i+1)^{th}$ entry of $\beta^*_{\mathbb{X}}$.

Proof. Use Proposition 4.2 and the identity $H_{\mathbb{X}}(i,j) = \sum_{(h,k)\leq(i,j)} c_{h,k}$ to compute α^*_{j+1}:
$$\begin{aligned} \alpha^*_{j+1} &= H_{\mathbb{X}}(t-1,j) - H_{\mathbb{X}}(t-1,j-1) \\ &= \sum_{(h,k)\leq(t-1,j)} c_{h,k} - \sum_{(h,k)\leq(t-1,j-1)} c_{h,k} = \sum_{h\leq t-1=|\pi_1(\mathbb{X})|-1} c_{h,j}. \end{aligned}$$
The proof for the second statement is the same. □

Lemma 4.6. *Let $\mathbb{X} \subseteq \mathbb{P}^1 \times \mathbb{P}^1$ be a finite set of points, and suppose that $\alpha^*_{\mathbb{X}} = \beta_{\mathbb{X}}$. Let P be a point of $\pi_1(\mathbb{X})$ such that $|\pi_1^{-1}(P)| = \alpha_1$. Set $\mathbb{X}_P := \pi_1^{-1}(P)$. Then $\pi_2(\mathbb{X}_P) = \pi_2(\mathbb{X})$.*

Proof. Since $\mathbb{X}_P \subseteq \mathbb{X}$, we have $\pi_2(\mathbb{X}_P) \subseteq \pi_2(\mathbb{X})$. Now, by our choice of P, $|\pi_2(\mathbb{X}_P)| = \alpha_1$. But since $|\pi_2(\mathbb{X})| = |\beta_{\mathbb{X}}|$ and $\alpha^*_{\mathbb{X}} = \beta_{\mathbb{X}}$, from Lemma 4.1 we have $|\pi_2(\mathbb{X})| = |\beta_{\mathbb{X}}| = \alpha_1 = |\pi_2(\mathbb{X}_P)|$, and hence $\pi_2(\mathbb{X}_P) = \pi_2(\mathbb{X})$. □

Proposition 4.7. *Suppose that* $\mathbb{X}$ *is a set of* $s = tr$ *points in* $\mathbb{P}^1 \times \mathbb{P}^1$ *such that* $\alpha_{\mathbb{X}} = (r, \ldots, r)$ *(*t *times) and* $\beta_{\mathbb{X}} = (t, \ldots, t)$ *(*r *times). Then* $\mathbb{X}$ *is a complete intersection, and the graded minimal free resolution of* $I_{\mathbb{X}}$ *is given by*

$$0 \longrightarrow R(-t,-r) \longrightarrow R(-t,0) \oplus R(0,-r) \longrightarrow I_{\mathbb{X}} \longrightarrow 0.$$

Proof. Because $|\alpha_{\mathbb{X}}| = t$ and $|\beta_{\mathbb{X}}| = r$, $\pi_1(\mathbb{X}) = \{P_1, \ldots, P_t\}$ and $\pi_2(\mathbb{X}) = \{Q_1, \ldots, Q_r\}$ where $P_i, Q_j \in \mathbb{P}^1$. Since $|\mathbb{X}| = tr$, $\mathbb{X} = \{P_i \times Q_j \mid 1 \leq i \leq t, 1 \leq j \leq r\}$. Hence, if $I_{P_i \times Q_j} = (L_{P_i}, L_{Q_j})$ is the ideal associated to the point $P_i \times Q_j$, then the defining ideal of $\mathbb{X}$ is

$$I_{\mathbb{X}} = \bigcap_{i,j} (L_{P_i}, L_{Q_j}) = (L_{P_1} L_{P_2} \cdots L_{P_t}, L_{Q_1} L_{Q_2} \cdots L_{Q_r}) .$$

Since $\deg L_{P_1} L_{P_2} \cdots L_{P_t} = (t, 0)$ and $\deg L_{Q_1} L_{Q_2} \cdots L_{Q_r} = (0, r)$, the two generators form a regular sequence on R, and hence, $\mathbb{X}$ is a complete intersection. The graded minimal free resolution is then given by the *Koszul resolution*, taking into consideration that $I_{\mathbb{X}}$ is bigraded. □

We now come to the main result of this section.

Theorem 4.8. *Let* $\mathbb{X}$ *be a finite set of points in* $\mathbb{P}^1 \times \mathbb{P}^1$ *with Hilbert function* $H_{\mathbb{X}}$*. Then the following are equivalent:*

- (i) $\mathbb{X}$ *is ACM.*
- (ii) $\Delta H_{\mathbb{X}}$ *is the Hilbert function of an* $\mathbb{N}^2$*-graded artinian quotient of* $\mathbf{k}[x_1, y_1]$.
- (iii) $\alpha^*_{\mathbb{X}} = \beta_{\mathbb{X}}$.

Proof. The implication $(i) \Rightarrow (ii)$ is Corollary 3.5. So, suppose that (ii) holds. Because $\Delta H_{\mathbb{X}}$ is the Hilbert function of an $\mathbb{N}^2$-graded artinian quotient of $\mathbf{k}[x_1, y_1]$, Corollary 3.14, Remark 3.15, and the matrix (1) give

$\Delta H_{\mathbb{X}} =$



where $t = |\pi_1(\mathbb{X})|$ and $r = |\pi_2(\mathbb{X})|$. We have written $\Delta H_{\mathbb{X}}$ as an infinite matrix whose indexing starts from zero rather than one.

By Corollary 4.5 the number of 1's in the $(i-1)^{th}$ row of $\Delta H_{\mathbb{X}}$ for each integer $1 \leq i \leq t$ is simply the i^{th} coordinate of $\beta^*_{\mathbb{X}}$. Similarly, the number of ones in the $(j-1)^{th}$ column of $\Delta H_{\mathbb{X}}$ for each integer $1 \leq j \leq r$ is the j^{th} coordinate of $\alpha^*_{\mathbb{X}}$. Now $\Delta H_{\mathbb{X}}$ can be identified with the Ferrers diagram of $\beta^*_{\mathbb{X}}$ by associating to each 1 in $\Delta H_{\mathbb{X}}$ a dot in the Ferrers diagram in the natural way, i.e.,



By using the Ferrers diagram and Corollary 4.5, it is now straightforward to calculate that the conjugate of $\beta_{\mathbb{X}}^*$ is $(\beta_{\mathbb{X}}^*)^* = \beta_{\mathbb{X}} = \alpha_{\mathbb{X}}^*$, and so (iii) holds.

To demonstrate that (iii) implies (i), we proceed by induction on the tuple $(|\pi_1(\mathbb{X})|, |\mathbb{X}|)$. For any positive integer s, if $(|\pi_1(\mathbb{X})|, |\mathbb{X}|) = (1, s)$, then $\alpha_{\mathbb{X}} = (s)$ and $\beta_{\mathbb{X}} = (1, \ldots, 1)$ (s times), and so $\alpha_{\mathbb{X}}^* = \beta_{\mathbb{X}}$. Then by Proposition 4.7 $\mathbb{X}$ is a complete intersection, and hence, ACM.

So, suppose that $(|\pi(\mathbb{X})), |\mathbb{X}|) = (t, s)$ and that result holds true for all $\mathbb{Y} \subseteq \mathbb{P}^1 \times \mathbb{P}^1$ with $\alpha_{\mathbb{Y}}^* = \beta_{\mathbb{Y}}$ and $(t, s) >_{lex} (|\pi_1(\mathbb{Y})|, |\mathbb{Y}|)$ where $>_{lex}$ is the lexicographical ordering on $\mathbb{N}^2$.

Suppose that P_1 (after a possible relabeling) is the element of $\pi_1(\mathbb{X})$ such that $|\pi_1^{-1}(P_1)| = \alpha_1$. Let L_{P_1} be the form of degree $(1, 0)$ that vanishes at P_1. By abusing notation, we also let L_{P_1} denote the $(1, 0)$-line in $\mathbb{P}^1 \times \mathbb{P}^1$ defined by L_{P_1}.

Set $\mathbb{X}_{P_1} := \mathbb{X} \cap L_{P_1} = \pi_1^{-1}(P_1)$ and $\mathbb{Z} := \mathbb{X} \backslash \mathbb{X}_{P_1}$. It follows that $\alpha_{\mathbb{Z}} = (\alpha_2, \ldots, \alpha_t)$ and $\beta_{\mathbb{Z}} = (\beta_1 - 1, \ldots, \beta_{\alpha_2} - 1)$. Now $(t, s) >_{lex} (|\pi_1(\mathbb{Z})|, |\mathbb{Z}|)$, and moreover, $\alpha_{\mathbb{Z}}^* = \beta_{\mathbb{Z}}$ by Lemma 4.1. Thus, by the induction hypothesis, $\mathbb{Z}$ is ACM.

Suppose that $\pi_2(\mathbb{X}) = \{Q_1, \ldots, Q_r\}$. Let L_{Q_i} be the degree $(0, 1)$ form that vanishes at $Q_i \in \pi_2(\mathbb{X})$ and set $F := L_{Q_1} L_{Q_2} \cdots L_{Q_r}$. Because $\alpha_{\mathbb{X}}^* = \beta_{\mathbb{X}}$, from Lemma 4.6 we have $\pi_2(\mathbb{X}_{P_1}) = \pi_2(\mathbb{X})$. So, $\mathbb{X}_{P_1} = \{P_1 \times Q_1, \ldots, P_1 \times Q_r\}$, and hence $I_{\mathbb{X}_{P_1}} = \bigcap_{i=1}^r (L_{P_1}, L_{Q_i}) = (L_{P_1}, F)$. Furthermore, if $P \times Q \in \mathbb{Z}$, then $Q \in \pi_2(\mathbb{Z}) \subseteq \pi_2(\mathbb{X})$, and thus $F(P \times Q) = 0$. Therefore $F \in I_{\mathbb{Z}}$. Because F is in $I_{\mathbb{Z}}$ and is also a generator of $I_{\mathbb{X}_{P_1}}$, we can show:

Claim. Let $I = L_{P_1} \cdot I_{\mathbb{Z}} + (F)$. Then $I = I_{\mathbb{X}}$.

Proof of the Claim. Since $I_{\mathbb{X}} = I_{\mathbb{Z} \cup \mathbb{X}_{P_1}} = I_{\mathbb{Z}} \cap I_{\mathbb{X}_{P_1}}$, we will show $I_{\mathbb{Z}} \cap I_{\mathbb{X}_{P_1}} = L_{P_1} \cdot I_{\mathbb{Z}} + (F)$. So, suppose that $G = L_{P_1} H_1 + H_2 F \in L_{P_1} \cdot I_{\mathbb{Z}} + (F)$ with $H_1 \in I_{\mathbb{Z}}$ and $H_2 \in R$. Because L_{P_1} and F are in $I_{\mathbb{X}_{P_1}}$, we have $G \in I_{\mathbb{X}_{P_1}}$. Since $H_1, F \in I_{\mathbb{Z}}$, $G \in I_{\mathbb{Z}}$. Thus $G \in I_{\mathbb{Z}} \cap I_{\mathbb{X}_{P_1}}$.

Conversely, let $G \in I_{\mathbb{Z}} \cap I_{\mathbb{X}_{P_1}}$. Since $G \in I_{\mathbb{X}_{P_1}}$, $G = L_{P_1} H_1 + F H_2$. We need to show that $H_1 \in I_{\mathbb{Z}}$. Because $G, F \in I_{\mathbb{Z}}$, we also have $L_{P_1} H_1 \in I_{\mathbb{Z}}$. But for every $P \times Q \in \mathbb{Z}$, $P \neq P_1$, and thus $L_{P_1}(P \times Q) \neq 0$. Hence $L_{P_1} H_1 \in I_{\mathbb{Z}}$ if and only if $H_1(P \times Q) = 0$ for every $P \times Q \in \mathbb{Z}$. □

We note that $\mathbb{X} \subseteq \mathbb{P}^1 \times \mathbb{P}^1$ is ACM if and only if the variety $\widetilde{\mathbb{X}} \subseteq \mathbb{P}^3$ defined by $I_{\mathbb{X}}$, considered as a homogeneous ideal of $R = \mathbf{k}[x_0, x_1, y_0, y_1]$, is ACM. As a variety of $\mathbb{P}^3$, $\widetilde{\mathbb{X}}$ is a curve since $\operatorname{K-dim} R/I_{\mathbb{X}} = 2$. Let $\widetilde{\mathbb{Z}}$ denote the curve of $\mathbb{P}^3$ defined by $I_{\mathbb{Z}}$, considered also as a homogeneous ideal of R. The claim implies that the curve $\widetilde{\mathbb{X}}$ is a basic double link of $\widetilde{\mathbb{Z}}$. Since the Cohen-Macaulay property is preserved under linkage (see [8, Theorem 3.2.3] and following remark), $\widetilde{\mathbb{X}}$ is an ACM curve of $\mathbb{P}^3$, or equivalently, $\mathbb{X}$ is an ACM set of points in $\mathbb{P}^1 \times \mathbb{P}^1$. □

Remark 4.9. Giuffrida, *et al.* [5, Theorem 4.1] demonstrated the equivalence of statements (i) and (ii) of Theorem 4.8 via different means. Our contribution is to show that the ACM sets of points are also characterized by $\alpha_{\mathbb{X}}$ and $\beta_{\mathbb{X}}$. This result has been extended in [6] to characterize ACM fat point schemes in $\mathbb{P}^1 \times \mathbb{P}^1$

Corollary 4.10. *Let $\mathbb{X}$ be a set of points in $\mathbb{P}^1 \times \mathbb{P}^1$ with $\alpha_{\mathbb{X}} = (\alpha_1, \ldots, \alpha_t)$, and $\pi_1(\mathbb{X}) = \{P_1, \ldots, P_t\}$. Suppose (after a possible relabeling) that $|\pi_1^{-1}(P_i)| = \alpha_i$. Set*

$$\mathbb{X}_i := \mathbb{X} \backslash \left\{\pi_1^{-1}(P_1) \cup \cdots \cup \pi_1^{-1}(P_i)\right\} \quad \text{for } 0 \leq i \leq t-1,$$

where $\mathbb{X}_0 := \mathbb{X}$. If $\mathbb{X}$ is ACM, then, for each integer $0 \leq i \leq t-1$, $\mathbb{X}_i$ is ACM with $\alpha_{\mathbb{X}_i} = (\alpha_{i+1}, \alpha_{i+2}, \ldots, \alpha_t)$.

Proof. It is sufficient to show that for each $i = 0, \ldots, t-2$, if $\mathbb{X}_i$ is ACM, then $\mathbb{X}_{i+1}$ is ACM. Since $\mathbb{X}_{i+1} = \mathbb{X}_i \backslash \{\pi_1^{-1}(P_{i+1})\}$, $\mathbb{X}_{i+1}$ is constructed from $\mathbb{X}_i$ by removing the α_{i+1} points of $\mathbb{X}_i$ which have P_{i+1} as its first coordinate. The tuple $\beta_{\mathbb{X}_{i+1}}$ is constructed from $\beta_{\mathbb{X}_i}$ by subtracting 1 from α_{i+1} coordinates in $\beta_{\mathbb{X}_i}$. But because $\alpha^*_{\mathbb{X}_i} = \beta_{\mathbb{X}_i}$, we have $|\beta_{\mathbb{X}_i}| = \alpha_{i+1}$, and thus $\beta_{\mathbb{X}_{i+1}} = (\beta_1 - 1, \ldots, \beta_{\alpha_{i+1}} - 1) = (\beta_1 - 1, \ldots, \beta_{\alpha_{i+2}} - 1)$. But by Lemma 4.1, $\alpha^*_{\mathbb{X}_{i+1}} = \beta_{\mathbb{X}_{i+1}}$, and hence, $\mathbb{X}_{i+1}$ is ACM by Theorem 4.8. □

5. The Betti numbers of ACM sets of points in $\mathbb{P}^1 \times \mathbb{P}^1$

If $\mathbb{X}$ is a set of s points in $\mathbb{P}^1$, then it is well known that the graded Betti numbers, and consequently, the Hilbert function of $\mathbb{X}$, can be determined solely from $|\mathbb{X}| = s$. If we restrict to ACM sets of points in $\mathbb{P}^1 \times \mathbb{P}^1$, we can extend this result to show that the graded Betti numbers in the minimal free resolution of $\mathbb{X}$ (and hence, $H_{\mathbb{X}}$) can be computed directly from the tuples $\alpha_{\mathbb{X}}$ and $\beta_{\mathbb{X}}$ introduced in the previous section which numerically describe $\mathbb{X}$.

We require the following notation to describe the minimal free resolution. Suppose that $\mathbb{X} \subseteq \mathbb{P}^1 \times \mathbb{P}^1$ is a set of points with $\alpha_{\mathbb{X}} = (\alpha_1, \ldots, \alpha_t)$. Define the following sets:

$$\begin{aligned} C_{\mathbb{X}} &:= \{(t,0),(0,\alpha_1)\} \cup \{(i-1,\alpha_i) \mid \alpha_i - \alpha_{i-1} < 0\}, \\ V_{\mathbb{X}} &:= \{(t,\alpha_t)\} \cup \{(i-1,\alpha_{i-1}) \mid \alpha_i - \alpha_{i-1} < 0\}. \end{aligned}$$

We take $\alpha_{-1} = 0$. With this notation, we have

Theorem 5.1. *Suppose that $\mathbb{X}$ is an ACM set of points in $\mathbb{P}^1 \times \mathbb{P}^1$. Let $C_{\mathbb{X}}$ and $V_{\mathbb{X}}$ be constructed from $\alpha_{\mathbb{X}}$ as above. Then the graded minimal free resolution of $I_{\mathbb{X}}$ is given by*

$$0 \longrightarrow \bigoplus_{(v_1,v_2) \in V_{\mathbb{X}}} R(-v_1,-v_2) \longrightarrow \bigoplus_{(c_1,c_2) \in C_{\mathbb{X}}} R(-c_1,-c_2) \longrightarrow I_{\mathbb{X}} \longrightarrow 0.$$

Proof. We proceed by induction on the tuple $(|\pi_1(\mathbb{X})|, |\mathbb{X}|)$. If s is any integer, and $(|\pi_1(\mathbb{X}), |\mathbb{X}|) = (1, s)$, then $\alpha_{\mathbb{X}} = (s)$ and $\beta_{\mathbb{X}} = (1, \ldots, 1)$ (s times). The conclusion follows from Proposition 4.7 since $C_{\mathbb{X}} = \{(1,0),(0,s)\}$ and $V_{\mathbb{X}} = \{(1,s)\}$.

So, suppose $(|\pi_1(\mathbb{X}), |\mathbb{X}|) = (t, s)$ with $t > 1$. Then $\alpha_{\mathbb{X}} = (\underbrace{\alpha_1, \ldots, \alpha_1}_{l}, \alpha_{l+1}, \ldots, \alpha_t)$, i.e., $\alpha_{l+1} < \alpha_1$, but $\alpha_l = \alpha_1$. If $l = t$, then $\mathbb{X}$ is a complete intersection and the resolution is given by Proposition 4.7. The conclusion now follows because $C_{\mathbb{X}} = \{(l,0),(0,\alpha_1)\}$ and $V_{\mathbb{X}} = \{(l,\alpha_1)\}$.

If $l < t$, let $P_1, \ldots, P_l$ be the l points of $\pi_1(\mathbb{X})$ that have $|\pi_1^{-1}(P_i)| = \alpha_1$. Set $\mathbb{Y} = \pi_1^{-1}(P_1) \cup \cdots \cup \pi_1^{-1}(P_l)$. Because $\mathbb{X}$ is ACM, $\alpha_1 = |\beta_{\mathbb{X}}|$, and hence, $\mathbb{Y} = \{P_i \times Q_j \mid 1 \leq i \leq l,\ Q_j \in \pi_2(\mathbb{X})\}$. So, $\alpha_{\mathbb{Y}} = (\alpha_1, \ldots, \alpha_1)$ and $\beta_{\mathbb{Y}} = (l, \ldots, l)$, and thus $\mathbb{Y}$ is a complete intersection. In fact,

$I_{\mathbb{Y}} = (L_{P_1}\cdots L_{P_l}, L_{Q_1}\cdots L_{Q_{\alpha_1}})$ where L_{P_i} is the form of degree $(1,0)$ that vanishes at all the points of $\mathbb{P}^1\times\mathbb{P}^1$ which have P_i as their first coordinate, and L_{Q_i} is the form of degree $(0,1)$ that vanishes at all points $P\times Q\in\mathbb{P}^1\times\mathbb{P}^1$ such that $Q=Q_i$.

Let $F := L_{P_1}\cdots L_{P_l}$ and $G := L_{Q_1}\cdots L_{Q_{\alpha_1}}$. By Proposition 4.7 the resolution of $I_{\mathbb{Y}}$ is

$$0 \longrightarrow R(-l,-\alpha_1) \xrightarrow{\phi_2} R(-l,0)\oplus R(0,-\alpha_1) \xrightarrow{\phi_1} I_{\mathbb{Y}} \longrightarrow 0$$

where $\phi_1 = [F\ G]$ and $\phi_2 = \begin{bmatrix} G \\ -F\end{bmatrix}$. Let $\mathbb{Z} := \mathbb{X}\backslash\mathbb{Y}$. Since $\pi_2(\mathbb{Z})\subseteq\pi_2(\mathbb{X})$, it follows that $G = L_{Q_1}\cdots L_{Q_{\alpha_1}}\in I_{\mathbb{Z}}$. Hence, $\operatorname{im}\phi_2\subseteq I_{\mathbb{Z}}(-l,0)\oplus R(0,-\alpha_1)$.

Claim. $I_{\mathbb{X}} = F\cdot I_{\mathbb{Z}} + (G)$

Proof of the Claim. By construction, $\mathbb{X} = \mathbb{Z}\cup\mathbb{Y}$. Hence, we want to show that $I_{\mathbb{Z}}\cap I_{\mathbb{Y}} = F\cdot I_{\mathbb{Z}} + (G)$. The proof now follows as in the proof of the Claim in Theorem 4.8. □

From the above resolution for $I_{\mathbb{Y}}$, the claim, and the fact that $\operatorname{im}\phi_2\subseteq I_{\mathbb{Z}}(-l,0)\oplus R(0,-\alpha_1)$, we have the following short exact sequence of R-modules

$$0 \longrightarrow R(-l,-\alpha_1) \xrightarrow{\phi_2} I_{\mathbb{Z}}(-l,0)\oplus R(0,-\alpha_1) \xrightarrow{\phi_1} I_{\mathbb{X}} = F\cdot I_{\mathbb{Z}} + (G) \longrightarrow 0$$

where ϕ_1 and ϕ_2 are as above.

By Corollary 4.10 the set $\mathbb{Z}$ is ACM with $\alpha_{\mathbb{Z}} = (\alpha_{l+1},\ldots,\alpha_t)$. Therefore, the induction hypothesis holds for $\mathbb{Z}$. With the above short exact sequence, we can use the *mapping cone construction* (see Section 1.5 of [14]) to construct a resolution for $I_{\mathbb{X}}$. In particular, we get

$$0 \longrightarrow \left[\bigoplus_{(v_1,v_2)\in V_{\mathbb{Z}}} R(-(v_1+l),-v_2)\right]\oplus R(-l,-\alpha_1) \longrightarrow$$

$$\left[\bigoplus_{(c_1,c_2)\in C_{\mathbb{Z}}} R(-(c_1+l),-c_2)\right]\oplus R(0,\alpha_1) \longrightarrow I_{\mathbb{X}} \longrightarrow 0.$$

Since the resolution has length 2, and because $\mathbb{X}$ is ACM, the resolution of $I_{\mathbb{X}}$ cannot be made shorter by the Auslander-Buchsbaum formula (cf. [14, Theorem 4.4.15]).

To show that this resolution is minimal, it is enough to show that no tuple in the set $\{(c_1+l,c_2)\mid (c_1,c_2)\in C_{\mathbb{Z}}\}\cup\{(0,\alpha_1)\}$ is in the set $\{(v_1+l,v_2)\mid (v_1,v_2)\in V_{\mathbb{Z}}\}\cup\{(l,\alpha_1)\}$. By the induction hypothesis, we can assume that no $(c_1,c_2)\in C_{\mathbb{Z}}$ is in $V_{\mathbb{Z}}$, and hence, if $(c_1+l,c_2)\in\{(c_1+l,c_2)\mid (c_1,c_2)\in C_{\mathbb{Z}}\}$, then (c_1+l,c_2) is not in $\{(v_1+l,v_2)\mid (v_1,v_2)\in V_{\mathbb{Z}}\}$ If $(c_1+l,c_2) = (l,\alpha_1)$ for some $(c_1,c_2)\in C_{\mathbb{Z}}$, then this implies that $(0,\alpha_1)$. But this contradictions the induction hypothesis. Similarly, if $(0,\alpha_1)\in\{(v_1+l,v_2)\mid (v_1,v_2)\in V_{\mathbb{Z}}\}$, this implies $(-l,\alpha_1)\in V_{\mathbb{Z}}$, which is again a contradiction of the induction hypothesis. So the resolution above is minimal.

To complete the proof we only need to verify that

(*i*) $C_{\mathbb{X}} = \{(c_1+l,c_2)\mid (c_1,c_2)\in C_{\mathbb{Z}}\}\cup\{(0,\alpha_1)\}$.
(*ii*) $V_{\mathbb{X}} = \{(v_1+l,v_2)\mid (v_1,v_2)\in V_{\mathbb{Z}}\}\cup\{(l,\alpha_1)\}$.

Because the verification of these statements is tedious, but elementary, we omit the details. □

Remark 5.2. It was shown in [5, Theorem 4.1] that the graded Betti numbers for an ACM set of points $\mathbb{X}\subseteq\mathbb{P}^1\times\mathbb{P}^1$ could be determined via the first difference function $\Delta H_{\mathbb{X}}$, i.e.,



An element of $C_{\mathbb{X}}$, which [5] called a *corner* of $\Delta H_{\mathbb{X}}$, corresponds to a tuple (i,j) that is either $(t,0),(0,\alpha_1) = (0,r)$, or has the property that $\Delta H_{\mathbb{X}}(i,j) = 0$, but $\Delta H_{\mathbb{X}}(i-1,j) = \Delta H_{\mathbb{X}}(i,j-1) = 1$. We have labeled the corners of $\Delta H_{\mathbb{X}}$ with a c in the above diagram. An element of $V_{\mathbb{X}}$ is a *vertex*. A tuple (i,j) is called a vertex if $\Delta H_{\mathbb{X}}(i,j) = \Delta H_{\mathbb{X}}(i-1,j) = \Delta H_{\mathbb{X}}(i,j-1) = 0$, but $\Delta H_{\mathbb{X}}(i-1,j-1) = 1$. We have labeled the vertices of $\Delta H_{\mathbb{X}}$ with a v in the above diagram. Besides giving a new proof for the resolution of an ACM set of points in $\mathbb{P}^1 \times \mathbb{P}^1$, we have shown that the graded Betti numbers can be computed directly from the tuple $\alpha_{\mathbb{X}}$.

Using the resolution as given in Theorem 5.1 we can compute $H_{\mathbb{X}}$ directly from $\alpha_{\mathbb{X}}$ for any ACM set of points $\mathbb{X} \subseteq \mathbb{P}^1 \times \mathbb{P}^1$. Formally

Corollary 5.3. *Let $\mathbb{X}$ be an ACM set of points in $\mathbb{P}^1 \times \mathbb{P}^1$ with $\alpha_{\mathbb{X}} = (\alpha_1, \ldots \alpha_t)$. Then*

$$H_{\mathbb{X}} = \begin{bmatrix} 1 & 2 & \cdots & \alpha_1 - 1 & \alpha_1 & \alpha_1 & \cdots \\ 1 & 2 & \cdots & \alpha_1 - 1 & \alpha_1 & \alpha_1 & \cdots \\ \vdots & \vdots & & \vdots & \vdots & \vdots & \ddots \end{bmatrix} + \begin{bmatrix} 0 & 0 & \cdots & 0 & 0 & 0 & \cdots \\ 1 & 2 & \cdots & \alpha_2 - 1 & \alpha_2 & \alpha_2 & \cdots \\ 1 & 2 & \cdots & \alpha_2 - 1 & \alpha_2 & \alpha_2 & \cdots \\ \vdots & \vdots & & \vdots & \vdots & \vdots & \ddots \end{bmatrix} +$$

$$\begin{bmatrix} 0 & 0 & \cdots & 0 & 0 & 0 & \cdots \\ 0 & 0 & \cdots & 0 & 0 & 0 & \cdots \\ 1 & 2 & \cdots & \alpha_3 - 1 & \alpha_3 & \alpha_3 & \cdots \\ 1 & 2 & \cdots & \alpha_3 - 1 & \alpha_3 & \alpha_3 & \cdots \\ \vdots & \vdots & & \vdots & \vdots & \vdots & \ddots \end{bmatrix} + \cdots + \begin{bmatrix} 0 & 0 & \cdots & 0 & 0 & 0 & \cdots \\ \vdots & \vdots & & \vdots & \vdots & \vdots & \\ 0 & 0 & \cdots & 0 & 0 & 0 & \cdots \\ 1 & 2 & \cdots & \alpha_t - 1 & \alpha_t & \alpha_t & \cdots \\ 1 & 2 & \cdots & \alpha_t - 1 & \alpha_t & \alpha_t & \cdots \\ \vdots & \vdots & & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

ACKNOWLEDGMENTS

Some of the results in this paper were part of [12]. The computer program CoCoA [2] was invaluable in providing and checking examples in the early stages. I would like to thank A. Ragusa, L. Roberts, D. Wehlau, and the audience of the Curve Seminar for their comments and suggestions. I would also like to thank the Università di Genova, where part of this work was completed. I would especially like to thank my supervisor Tony Geramita for introducing me to this problem, and for his encouragement and help.

REFERENCES

[1] W. Bruns, J. Herzog (1998), *Cohen-Macaulay rings* (Revised Edition). Cambridge University Press, New York.

[2] A. Capani, G. Niesi, L. Robbiano, `CoCoA`, a system for doing Computations in Commutative Algebra, Available via anonymous ftp from: `cocoa.dima.unige.it`

[3] A.V. Geramita, D. Gregory, L. Roberts (1986), Monomial ideals and points in projective space. JPAA **40**, 33–62.

[4] A.V. Geramita, P. Maroscia, L. Roberts (1983), The Hilbert function of a reduced k-algebra. J. London Math. Soc. **28**, 443–452.

[5] S. Giuffrida, R. Maggioni, A. Ragusa (1992), On the postulation of 0-dimensional subschemes on a smooth quadric. Pacific J. Math. **155**, 251–282.

[6] E. Guardo, A. Van Tuyl (2002) Fat Points in $\mathbb{P}^1 \times \mathbb{P}^1$ and their Hilbert functions. To appear Can. J. Math.

[7] F.S. Macaulay (1927), Some properties of enumeration in the theory of modular systems. Proc. London Math. Soc. **26**, 531–555.

[8] J. Migliore (1998), *Introduction to Liaison Theory and Deficiency Modules.* Birkhäuser: Boston.

[9] J. Migliore, U. Nagel (2000), Lifting Monomial Ideals. Comm. Algebra **28**, 5679–5701.

[10] P. Roberts, *Multiplicites and Chern Classes in Local Algebra.* Cambridge Tracts in Mathematics, 133. Cambridge University Press, Cambridge, 1998.

[11] R. Stanley (1978), Hilbert functions of graded algebras. Adv. in Math. **28**, 57-83.

[12] A. Van Tuyl (2001), Sets of Points in Multi-Projective Spaces and their Hilbert Function, Ph.D. Thesis, Queen's University.

[13] A. Van Tuyl (2002), The border of the Hilbert function of a set of points in $\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$. JPAA **176**, 223-247.

[14] C. Weibel (1994), *An introduction to homologocial algebra.* Cambridge University Press: New York.

Department of Mathematical Sciences, Lakehead University, Thunder Bay, ON P7B 5E1, Canada

E-mail address: `avantuyl@sleet.lakeheadu.ca`